

Membrane paradigm in the Einstein-dilaton theory

Chanyong Park^{a*}

^a *Center for Quantum Spacetime (CQUeST), Sogang University, Seoul 121-742, Korea*

ABSTRACT

We study a general D -dimensional Schwarzschild-type black brane solution in the Einstein-dilaton theory and investigate its thermodynamics. Using the membrane paradigm we calculate the several hydrodynamic transport coefficients and then compare them with the results obtained by the Kubo formula in the 4-dimensional case. By calculating the momentum diffusion constant of the shear modes, we calculate the sum of the energy density and pressure, which is consistent with the result of the black brane thermodynamics.

*e-mail : cyong21@sogang.ac.kr

Contents

1	Introduction	1
2	<i>D</i>-dimensional black brane in the Einstein-dilaton theory	2
3	Membrane paradigm	5
3.1	DC conductivity	5
3.2	Charge diffusion constant	7
3.3	Shear modes	9
4	Discussion	11

1 Introduction

For the last decade, the AdS/CFT correspondence [1, 2, 3] has been one of the very interesting and fascinating subjects for understanding the strongly interacting quantum field theory (QFT). Applying the AdS/CFT correspondence to QCD or the condensed matter system provided many interesting physical results like the phase structures [4], the ratio between the shear viscosity and entropy [5, 6, 7, 8, 9, 10, 11], holographic superconductor [12]-[22] and strange metallic behavior [23], etc. In this paper, we will investigate the hydrodynamics for the relativistic nonconformal field theory dual to the Einstein-dilaton gravity by the generalized concept of the AdS/CFT correspondence, the so-called gauge/gravity duality.

Recently, after assuming the gauge/gravity duality it was shown that the dual field theory of the Einstein-dilaton gravity can be described by the relativistic nonconformal theory [24], in which the DC conductivity and the charge diffusion constant were calculated by using the Kubo formula [5, 6, 24]. The DC conductivity with the running dilaton coupling shows the unexpected behavior which may describe electrolyte. In this paper, following the membrane paradigm, we will generalize this gauge/gravity duality to the general D -dimensional Einstein-dilaton theory and furthermore apply it to the shear modes. The shear modes are important to check this gauge/gravity duality because they contain information for $\epsilon + P$ in the thermal equilibrium, where ϵ and P implies the energy density and pressure respectively. Independently, the same result can be also determined by the thermodynamics of the black brane. If the gauge/gravity duality is working in the Einstein-dilaton theory, those two results should be consistent. Notice that the black brane thermodynamics does not depend on the details of the dual QFT, whereas the hydrodynamic calculation needs techniques or informations of the quantum field theory. As a result, the calculation of $\epsilon + P$ becomes a nontrivial check of the gauge/gravity duality in the Einstein-dilaton theory, although the exact map between the Einstein-dilaton gravity and its dual QFT is not clear.

The rest parts are following: In Sec.2, we consider an Einstein-dilaton gravity with a Liouville-type dilaton potential and find a black brane solution with arbitrary constant η in the general dimension D . By investigating the thermodynamics of this black brane, we show that according to the gauge/gravity duality the dual field theory is described by the relativistic non-conformal field theory at finite temperature. Sec.3 is devoted to the computation of several transport coefficients of the dual non-conformal theory. Using the membrane paradigm, we study the DC conductivity and charge diffusion constant, which are consistent with those obtained by the Kubo formula in the 4-dimension [24]. Moreover, we also investigate the shear viscosity and momentum diffusion constant, which are important quantities because they involves informations for the dual non-conformal field theory, as we mentioned previously. We show that the thermodynamic property $\epsilon + P$ can be also reproduced from the hydrodynamics of the dual field theory. In Sec. 4, we finish this work with some concluding remarks.

2 D -dimensional black brane in the Einstein-dilaton theory

We consider a general D -dimensional Einstein-dilaton theory, which is the generalization of the 4-dimensional warped geometry [24] and provides a hyperscaling violating solution [25]-[35]. The action is

$$S = \frac{1}{16\pi G} \int d^D x \sqrt{-g} [R - 2(\partial\phi)^2 - V(\phi)], \quad (1)$$

with a Liouville-type potential

$$V(\phi) = 2\Lambda e^{\eta\phi}, \quad (2)$$

where Λ is a negative cosmological constant ($\Lambda < 0$) and η is an arbitrary constant representing the non-conformality of the dual theory respectively. For convenience, we set $G = 1$ from now on. The Einstein equation and equation of motion for dilaton are given by

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \frac{1}{2}g_{\mu\nu}V(\phi) = 2\partial_\mu\phi\partial_\nu\phi - g_{\mu\nu}(\partial\phi)^2, \quad (3)$$

$$\frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}g^{\mu\nu}\partial_\nu\phi) = \frac{1}{4}\frac{\partial V(\phi)}{\partial\phi}. \quad (4)$$

To solve these equations, we take the following general metric ansatz for a black brane

$$\begin{aligned} ds^2 &\equiv -g_{tt} dt^2 + g_{rr} dr^2 + g_{ij} dx^i dx^j \\ &= -a(r)^2 f(r) dt^2 + \frac{dr^2}{a(r)^2 f(r)} + b(r)^2 \delta_{ij} dx^i dx^j, \end{aligned} \quad (5)$$

with

$$\begin{aligned} \phi(r) &= \phi_0 - k_0 \log r, \\ a(r) &= a_0 r^{a_1}, \\ b(r) &= b_0 r^{b_1}, \\ f(r) &= 1 - \frac{m}{r^c}, \end{aligned} \quad (6)$$

where i and j represent the spatial coordinates of the boundary space and m is the black brane mass which appears as an integration constant. Notice that we can set, without loss of generality, $b_0 = 1$ by using the rescaling of spatial coordinates x^i and absorb ϕ_0 into the effective cosmological constant $\Lambda_{eff} = \Lambda e^{\eta\phi_0}$. The solutions satisfying all equations are given by

$$\begin{aligned} k_0 &= \frac{2(D-2)\eta}{8 + (D-2)\eta^2}, \\ a_1 = b_1 &= \frac{8}{8 + (D-2)\eta^2}, \\ a_0 &= \frac{(8 + (D-2)\eta^2) \sqrt{-\Lambda_{eff}}}{2\sqrt{(D-2)[8(D-1) - (D-2)\eta^2]}}, \\ c &= \frac{8(D-1) - (D-2)\eta^2}{8 + (D-2)\eta^2}. \end{aligned} \quad (7)$$

This black brane solution is valid only for $\eta^2 < 8(D-1)/(D-2)$, which is called the Gubser bound [36, 37, 38, 39]. If not, a_0 becomes an imaginary number, which exchanges the roles of time and the radial coordinates. Notice that since the redefinition of time as $t \rightarrow t/a_0$ is also possible, we can set $a_0 = 1$. Here, we simply set

$$\Lambda_{eff} = -\frac{4(D-2)[8(D-1) - (D-2)\eta^2]}{[8 + (D-2)\eta^2]^2}, \quad (8)$$

which fix the combination of the cosmological constant and the integration constant ϕ_0 . As a result, the asymptotic form of the black brane metric in the Einstein-dilaton theory reduces to

$$ds^2 = \frac{1}{r^{2a_1}} dr^2 + r^{2a_1} (-dt^2 + dx^2 + dy^2). \quad (9)$$

In this case, the Poincare symmetry $ISO(1,2)$ of the hypersurface located at a fixed r becomes clear, so the dual field theory is described by a relativistic QFT. For $\eta = 0$, the background geometry reduces to the asymptotic AdS space with the effective cosmological constant Λ_{eff}

$$\Lambda_{eff} = -\frac{(D-1)(D-2)}{2}, \quad (10)$$

which is the cosmological constant of a general D -dimensional AdS space, as we expected.

For more understanding on the dual field theory, we need to study the thermodynamics of this black brane because it can be identified with that of the dual theory. From the metric in (5) together with (6) and (7), the event horizon r_h is located at

$$r_h = m^{-[8+(D-2)\eta^2]/[(D-2)\eta^2-8(D-1)]}. \quad (11)$$

Then, the Hawking temperature T_H defined by the surface gravity can be represented, in terms of the horizon, as

$$T_H = \frac{(-\Lambda_{eff})}{16\pi} \frac{8 + (D-2)\eta^2}{(D-2)} r_h^{[8-(D-2)\eta^2]/[8+(D-2)\eta^2]}. \quad (12)$$

and the Bekenstein-Hawking entropy S_{BH} becomes

$$S_{BH} = \frac{V_{D-2}}{4} r_h^{8(D-2)/[8+(D-2)\eta^2]}, \quad (13)$$

where V_{D-2} is the spatial volume of the boundary space. Usually, a black hole (or brane) system provides a well-defined analogous thermodynamic system, so the above black brane should satisfy the first thermodynamic law

$$0 = dE - T_H dS_{BH}. \quad (14)$$

From this relation we can determine the energy of the black brane geometry by rewriting the Hawking temperature in terms of the Bekenstein-Hawking entropy and then integrating it. In terms of the event horizon, the energy becomes

$$E = \frac{(-\Lambda_{eff})V_{D-2}}{8\pi} \frac{8 + (D-2)\eta^2}{8(D-1) - (D-2)\eta^2} r_h^{[8(D-1)-(D-2)\eta^2]/[8+(D-2)\eta^2]}, \quad (15)$$

and the related free energy is given by

$$F = -\frac{(-\Lambda_{eff})V_{D-2}}{64\pi} \frac{64 - (D-2)^2\eta^4}{(D-2)[8(D-1) - (D-2)\eta^2]} r_h^{[8(D-1)-(D-2)\eta^2]/[8+(D-2)\eta^2]}. \quad (16)$$

The pressure of the system can be easily obtained from the thermodynamic relation $P = -\partial F/\partial V_{D-2}$, so we can easily read off the equation of state parameter from the following relation $P = wE/V_2$

$$w = \frac{1}{D-2} - \frac{\eta^2}{8}, \quad (17)$$

which shows that the dual gauge theory with a general η is not conformal. For $\eta = 0$, the asymptotic geometry is the D -dimensional AdS space and the energy-momentum tensor of its dual field theory becomes traceless, as we expected. For the general η , since $0 \leq \eta^2 < 8(D-1)/(D-2)$, the equation of state parameter w can have the following values

$$-1 < \omega \leq \frac{1}{D-2}. \quad (18)$$

To check the thermodynamic stability of the dual field theory, we calculate the heat capacity in terms of Hawking temperature

$$C_V = \frac{2(D-2)V_{D-2}}{8 - (D-2)\eta^2} \left(\frac{16(D-2)\pi}{[8 + (D-2)\eta^2](-\Lambda_{eff})} \right)^{8(D-2)/[8-(D-2)\eta^2]} T_H^{8(D-2)/[8-(D-2)\eta^2]}. \quad (19)$$

Note that the heat capacity of the AdS black brane can be reproduced by setting $\eta = 0$ and it is always positive, which implies that the AdS black brane is thermodynamically stable. For $\eta^2 < 8/(D-2)$, the black brane has also a positive heat capacity, while its heat capacity is singular for $\eta^2 = 8/(D-2)$ and negative for $8/(D-2) < \eta^2 < 8(D-1)/(D-2)$. As a result, the Schwarzschild-type black brane of the Einstein-dilaton theory is thermodynamically stable only for $0 \leq \eta^2 < 8/(D-2)$ which implies that the dual field theory is generally described by a relativistic non-conformal quantum field theory and thermodynamically stable only for $0 < w \leq 1/(D-2)$. Notice that if taking $D = 4$, we can also reproduce all results in [24].

3 Membrane paradigm

Following the usual AdS/CFT correspondence, the dual operator of a bulk vector fluctuation A^i is a boundary current j^i . The one-point correlation function of j^i in the quantum field theory (QFT) can be defined by

$$\langle j^i(k_\mu) \rangle_{QFT} \equiv j^i(k_\mu) = \sigma^{ij}(k_\mu) F_{jt}, \quad (20)$$

where k_μ is the covariant momentum vector and $F_{jt} = E_j$ corresponds to the electric field in j -direction. In general, since σ^{ij} is a function of the frequency $k_t = \omega$ it corresponds to the AC conductivity which is also related to the retarded Green function

$$\sigma^{ij}(k_\mu) = -\frac{G_R^{ij}(k_\mu)}{i\omega}. \quad (21)$$

In terms of the canonical conjugate momentum Π^{ri} of a bulk vector field A_i , the retarded Green function can be also rewritten as

$$G_R^{ij}(k_\mu) = -\lim_{r \rightarrow \infty} \frac{\Pi^{ri}(r, k_\mu)}{A_j(r, k_\mu)}. \quad (22)$$

In the zero momentum limit $k_i = 0$ (i means the spatial directions), σ^{ij} can be represented as $\sigma^{ij} = \sigma_{AC}(\omega)\delta^{ij}$ due to the rotational symmetry. In this zero momentum limit, the DC conductivity can be obtained by taking the zero frequency limit

$$\sigma_{DC} = \lim_{\omega \rightarrow 0} \sigma_{AC}(\omega). \quad (23)$$

3.1 DC conductivity

Now, we consider a stretched horizon r_s which is a timelike surface at a fixed r just outside of the event horizon. Since no observer can hover at the event horizon, the stretched horizon acts as a cutoff for the bulk space outside the black brane. In the membrane paradigm, the dual field theory is defined on the membrane located at this stretched horizon instead of the asymptotic boundary. In other words, the bulk fluctuations are dual to some operators on the membrane. Then, the generating functional of those operators is represented by the surface term of the action on the membrane instead of the asymptotic boundary. In this section, we will study the hydrodynamic transport coefficients of the dual QFT by using a membrane paradigm [8, 9, 10] and compare them with the results of the Kubo formula [24]. Notice that as will be shown, the membrane paradigm is a very powerful method to obtain the general results in the general D -dimension with the general parameters.

Regard a bulk $U(1)$ gauge field with a r -dependent gauge coupling $g_D(r)^2$

$$S = - \int d^D x \sqrt{-g} \frac{1}{4g_D(r)^2} F_{MN} F^{MN}, \quad (24)$$

where $1/g_D(r)^2 = e^{\alpha\phi(r)}/g_0^2$ and g_0^2 is a constant. First, we concentrate on a DC conductivity on the membrane located at the stretched horizon. Since a DC conductivity is well-defined without the spatial momentum, we can take the zero momentum limit without loss of generality. In the $A_r = 0$ gauge, the Fourier mode of the vector field A_μ can be represented as

$$A_\mu(r, t) = \int \frac{d\omega}{2\pi} e^{-i\omega t} A_\mu(r, \omega). \quad (25)$$

If regarding r as a time coordinate, the canonical conjugate momentum Π^{ri} of the spatial component of A_i , which can be identified with the current j^i on the membrane, is given by

$$\Pi^{ri} \equiv j^i = -\frac{\sqrt{-g}}{g_D(r)^2} g^{rr} g^{ii} F_{ri}. \quad (26)$$

In the usual gauge/gravity duality, an incoming boundary condition at the horizon plays an important role to understanding the physical properties of the dual QFT, which is also true in the membrane paradigm. Since the horizon is a regular place to a free falling observer, the electromagnetic field detected by him must be also regular. This implies that near the horizon the bulk gauge fluctuation should be depend on r and t through their non-singular combination. In the Eddington-Finkelstein coordinate

$$u = dt + \sqrt{\frac{g_{rr}}{g_{tt}}} dr \quad \text{and} \quad v = dt - \sqrt{\frac{g_{rr}}{g_{tt}}} dr, \quad (27)$$

u and v describe the incoming and outgoing modes respectively. In this coordinate, the incoming boundary condition at the horizon implies that the electromagnetic field should be given by a function of u only, so $F_{vi} = \partial_v A_i = 0$. By using the chain rule, this incoming boundary condition can be rewritten as

$$F_{ri} = \sqrt{\frac{g_{rr}}{g_{tt}}} F_{ti}, \quad (28)$$

which implies that all fluctuations around the black brane must have only the incoming modes at least in the classical gravity theory. Then, the current on the membrane can be rewritten as

$$j^i = \frac{1}{g_D(r)^2} \sqrt{\frac{-g}{g_{tt}g_{rr}}} g^{ii} F_{it} \Big|_{r_s}. \quad (29)$$

Comparing it with (20), the conductivity of the membrane lying in the stretched horizon is given by

$$\sigma = \frac{1}{g_D(r)^2} \sqrt{\frac{-g}{g_{tt}g_{rr}}} g^{ii} \Big|_{r_s}. \quad (30)$$

The equations of motion and Bianchi identity $0 = \partial_{[r} F_{it]}$ become in terms of the current

$$\partial_r j^i = i\omega \frac{\sqrt{-g}}{g_D(r)^2} g^{tt} g^{ii} F_{it}, \quad (31)$$

$$\partial_r F_{it} = -i\omega \frac{g_D(r)^2}{\sqrt{-g}} g_{rr} g_{ii} j^i, \quad (32)$$

where we used (26) and (28). In the zero frequency limit, the current j^i and electric field F_{it} are independent of the radial coordinate, which implies that the DC conductivity in (30) actually holds for all r . If we put r_s into r_h and rewrite r_h in terms of the Hawking temperature (12), the DC conductivity in the general dimension D with the general parameters, α and η , reduces to

$$\sigma_{DC} = \frac{1}{g_0^2} \left(\frac{16(D-2)\pi}{[8 + (D-2)\eta^2](-\Lambda_{eff})} \right)^{\frac{8(D-4)-2(D-2)\eta\alpha}{8-(D-2)\eta^2}} T_H^{\frac{8(D-4)-2(D-2)\eta\alpha}{8-(D-2)\eta^2}}. \quad (33)$$

Especially, the DC conductivities, after taking special parameter values, become

$$\sigma_{DC} = \frac{1}{g_0^2} \quad \text{for } D = 4 \text{ and } \alpha = 0, \quad (34)$$

and

$$\sigma_{DC} = \frac{1}{g_0^2} \left(\frac{16\pi}{(4 + \eta^2)(-\Lambda_{eff})} \right)^{\frac{\eta^2}{4-\eta^2}} T_H^{\frac{\eta^2}{4-\eta^2}} \quad \text{for } D = 4 \text{ and } \alpha = -\eta/2, \quad (35)$$

which are coincident with the results in [24].

3.2 Charge diffusion constant

Consider a thermal equilibrium of a quantum field theory. If we disturb this system by turning on a small nonuniform perturbation of charge density varying along y -direction, the charge gradient generates a non-vanishing current, which eventually diffuses away back into thermal equilibrium. At the lowest order of the derivative expansion the diffusion process is governed by the following dispersion relation

$$\omega = -i\mathcal{D}k^2, \quad (36)$$

where $\mathcal{D}$ is the charge diffusion constant. In the linear response theory, the above appears as a pole of the retarded Green function of the longitudinal mode even in the absence of an external field. Following the gauge/gravity duality, this thermalization process can also be described by a dual gravity theory. Since the bulk black hole or brane solutions usually provide a well-defined thermodynamic concept to the dual field theory, the evolution of the black brane with corresponding fluctuation can explain the diffusion process of the dual field theory. Keeping it in mind, we will investigate the charge diffusion of QFT by studying the dual gravity theory.

If we take a momentum of the vector fluctuation to be along the y -direction $k = k_y$, the equations governing the vector fluctuations are divided into two parts: one is the longitudinal modes A_t and A_y and the other is a transverse mode A_x , where we still use the $A_r = 0$ gauge. Since the momentum plays an important role to determine the charge diffusive pole, we take the hydrodynamic regime, $\omega \sim k^2$, $\omega \ll T_H$, and $k \ll T_H$, instead of the zero momentum and zero frequency limit. As mentioned previously, the charge diffusion appears as a pole of the retarded Green function of the longitudinal modes only, so from now on we concentrate on the longitudinal modes. The longitudinal modes should

satisfy two dynamical equations as well as the conservation of j^μ and the Bianchi identity, which can be rewritten in terms of current j^μ in (29) as

$$0 = -\partial_r j^t - \frac{\sqrt{-g}}{g_D(r)^2} g^{tt} g^{yy} \partial_y F_{yt}, \quad (37)$$

$$0 = -\partial_r j^y - \frac{\sqrt{-g}}{g_D(r)^2} g^{tt} g^{yy} \partial_t F_{yt}, \quad (38)$$

$$0 = \partial_t j^t + \partial_y j^y, \quad (39)$$

$$0 = -\frac{g_{rr} g_{yy} g_D(r)^2}{\sqrt{-g}} \partial_t j^y - \frac{g_{rr} g_{yy} g_D(r)^2}{\sqrt{-g}} \partial_z j^t + \partial_r F_{yt}. \quad (40)$$

From these equations, we can derive a flow equation for the longitudinal conductivity $\sigma_L(r, k_\mu) \equiv j^y/F_{yt}$ [8]

$$\partial_r \sigma_L = i\omega \sqrt{\frac{g_{rr}}{g_{tt}}} \left[\frac{\sigma_L^2}{\Sigma_A(r)} \left(1 - \frac{k^2}{\omega^2} \frac{g^{yy}}{g^{tt}} \right) - \Sigma_A(r) \right], \quad (41)$$

with

$$\Sigma_A(r) = \frac{1}{g_D(r)^2} \sqrt{\frac{-g}{g_{tt} g_{rr}}} g^{yy}. \quad (42)$$

Similarly, using $g^{xx} = g^{yy}$, the flow equation for the transverse conductivity $\sigma_T \equiv j^x/F_{xt}$ is given by [8]

$$\partial_r \sigma_T = i\omega \sqrt{\frac{g_{rr}}{g_{tt}}} \left[\frac{\sigma_T^2}{\Sigma_A(r)} - \Sigma_A(r) \left(1 - \frac{k^2}{\omega^2} \frac{g^{yy}}{g^{tt}} \right) \right]. \quad (43)$$

In the zero frequency limit, regardless of the momentum, above two conductivities are independent of the position of the stretched horizon. Furthermore, imposing the regularity of the conductivities at the horizon they reduce the value of the DC conductivity in the previous section

$$\sigma_{L,T}(r_h, k_\mu) = \Sigma_A(r_h) = \sigma_{DC}, \quad (44)$$

which plays a role of the initial data for the conductivity flow. In the hydrodynamic limit ($\omega, k \ll T_H$), the flow equation of the longitudinal conductivity reduces to

$$\frac{\partial_r \sigma_L}{\sigma_L^2} = -i \frac{k^2}{\omega} \frac{g_{tt} g_{rr}}{\sqrt{-g}} g_D(r)^2. \quad (45)$$

According to the holographic renormalization, the radial position of the stretched horizon r_s can be identified with the energy scale of the dual field theory. Integrating (45) from r_h to r_s with the initial data in (44), the longitudinal conductivity at r_s becomes

$$\sigma_L(r_s) = \frac{i\omega \sigma_{DC}}{i\omega - \mathcal{D}(r_s) k^2}, \quad (46)$$

where the charge diffusion constant $\mathcal{D}(r_s)$ depending on the energy scale is given by

$$\mathcal{D}(r_s) = \sigma_{DC} \int_{r_h}^{r_s} dr \frac{g_{tt} g_{rr}}{\sqrt{-g}} g_D(r)^2. \quad (47)$$

This result together with (21) shows that the retarded Green function has the following form

$$G_R^{yy}(r_s, k_\mu) = \frac{\omega^2 \sigma_{DC}}{i\omega - \mathcal{D}(r_s)k^2}, \quad (48)$$

which is exactly the form obtained by the Kubo formula at the first order perturbation [24]. Using (21) and (23), we can reobtain the DC conductivity consistent with (30).

In the general D -dimensional Einstein-dilaton theory (5), the charge diffusion constant of the dual field theory becomes

$$\mathcal{D}(r_s) = \frac{(r_h^{-\delta} - r_s^{-\delta})}{\delta} r_h^\chi, \quad (49)$$

where

$$\begin{aligned} \delta &= \frac{8(D-3) - (D-2)(\eta + 2\alpha)\eta}{8 + (D-2)\eta^2}, \\ \chi &= \frac{8(D-4) - 2(D-2)\alpha\eta}{8 + (D-2)\eta^2}. \end{aligned} \quad (50)$$

For the positive δ , in other words $\alpha < \frac{8(D-3) - (D-2)\eta^2}{2(D-2)\eta}$, the charge diffusion constant of the dual field theory is well-defined at the limit of $r_s \rightarrow \infty$. Otherwise, it diverges at the infinity. For the well-defined hydrodynamic transport coefficients at the asymptotic boundary, we concentrate only on the positive δ from now on. Then, the general charge diffusion constant at the asymptotic boundary becomes

$$\mathcal{D}(\infty) = \frac{(-\Lambda_{eff})}{16\pi T_H} \frac{[8 + (D-2)\eta^2]^2}{(D-2)[8(D-3) - (D-2)(\eta + 2\alpha)\eta]}. \quad (51)$$

At the horizon $r_s \rightarrow r_h$, the charge diffusion constant becomes zero. From (49), we can easily see that the charge diffusion constant decreases monotonically from (51) along the RG flow. Especially, in the 4-dimension the charge diffusion constant at the asymptotic boundary ($r_s \rightarrow \infty$) reduces to

$$\mathcal{D} = \frac{(-\Lambda_{eff})}{16\pi T_H} \frac{(4 + \eta^2)^2}{4 - \eta^2} \quad \text{for } \alpha = 0, \quad (52)$$

and

$$\mathcal{D} = \frac{(-\Lambda_{eff})}{16\pi T_H} \frac{(4 + \eta^2)^2}{4} \quad \text{for } \alpha = -\frac{\eta}{2}, \quad (53)$$

which are exactly coincident with the results obtained by the Kubo formula [24].

3.3 Shear modes

As shown in [7, 8], the relevant equations for gravitational shear modes h_t^x and h_y^x , can be mapped to an electromagnetic problem. If we set $h_t^x = a_t$ and $h_y^x = a_y$, the action governing the shear mode fluctuations is reduced to

$$S = \frac{1}{16\pi} \int d^D x \sqrt{-g} g_{xx} F_{\alpha\beta} F^{\alpha\beta}, \quad (54)$$

where α and β imply the longitudinal direction t or y . Here, $F_{\alpha\beta}$ is the field strength of a_t and a_y , which in terms of metric fluctuations is given by $F_{\alpha\beta} = \partial_\alpha h_\beta^x - \partial_\beta h_\alpha^x$. This action for shear modes is exactly the standard Maxwell form with an effective coupling g_G^2

$$\frac{1}{g_G^2} = \frac{1}{16\pi} g_{xx}. \quad (55)$$

So we can immediately take over all results of the previous section. The corresponding DC conductivity σ_G at the horizon becomes

$$\sigma_G = \frac{1}{16\pi} \sqrt{\frac{-g}{g_{tt}g_{rr}}} \Big|_{r_h}. \quad (56)$$

Notice that the Bekenstein-Hawking entropy density s can be written as

$$s \equiv \frac{S_{BH}}{V_{D-2}} = \frac{1}{4} \sqrt{\frac{-g}{g_{tt}g_{rr}}} \Big|_{r_h}. \quad (57)$$

The real meaning of the above DC conductivity for the shear modes is the entropy density divided by 4π , which is the expected result in the Kubo formula and corresponds to the shear viscosity due to the celebrated universality of the ratio between the shear viscosity and the entropy [8]

$$\frac{\eta_s}{s} = \frac{1}{4\pi}. \quad (58)$$

Similarly, we can also easily evaluate the retarded Green function of the shear modes

$$G_R^{xy,xy} = \frac{\eta_s \omega^2}{i\omega - \mathcal{D}_s(r_s) k^2}, \quad (59)$$

where the momentum diffusion constant $\mathcal{D}_s$, by taking analogy with the charge diffusion constant, is

$$\begin{aligned} \mathcal{D}_s(r_s) &= 4s \int_{r_h}^{r_s} dr \frac{g_{tt} g_{rr}}{\sqrt{-g} g_{xx}} \\ &= \frac{8 + (D-2)\eta^2}{8(D-1) - (D-2)\eta^2} \left(\frac{1}{r_h^\gamma} - \frac{1}{r_s^\gamma} \right). \end{aligned} \quad (60)$$

with

$$\gamma = \frac{8 - (D-2)\eta^2}{8 + (D-2)\eta^2}. \quad (61)$$

Below the crossover value $\eta^2 < 8/(D-2)$ [37, 38, 39], the black brane is thermodynamically stable and γ is always positive. If we put r_s at the infinity, the momentum diffusion constant in the membrane paradigm reduces to

$$\mathcal{D}_s(r_s) = \frac{8 + (D-2)\eta^2}{8(D-1) - (D-2)\eta^2} \frac{1}{r_h^{[8-(D-2)\eta^2]/[8+(D-2)\eta^2]}}. \quad (62)$$

Note that the momentum diffusion constant at the infinite boundary is also related to the shear viscosity η_s in the hydrodynamics of QFT

$$\mathcal{D}_s(\infty) = \frac{\eta_s}{\epsilon + P}, \quad (63)$$

where ϵ and P are the energy density and pressure of the dual theory located at infinity respectively. In terms of the thermodynamic quantities in (13), (15), and (16), the momentum diffusion constant can be rewritten as

$$\frac{\eta_s}{\epsilon + P} = \frac{4(D-2)}{(8 + (D-2)\eta^2)(-\Lambda_{eff})} \frac{1}{r_h^{[8-(D-2)\eta^2]/[8+(D-2)\eta^2]}}. \quad (64)$$

Using the effective cosmological constant in (8), the momentum diffusion constants of the membrane paradigm and the hydrodynamics are consistent and finally reduces to

$$\mathcal{D}_s(\infty) = \frac{1}{4\pi T_H}, \quad (65)$$

which shows that the momentum diffusive pole decays rapidly at high temperature as we expected. In addition, it also shows that the momentum diffusion constant does not depend on the nonconformality for the dual field theory.

4 Discussion

In this paper, we have investigated the thermodynamic properties of the general D -dimensional Schwarzschild-type black brane in the Einstein-dilaton theory. Through the gauge/gravity duality, we found that the dual field theory for general η is described by a relativistic non-conformal quantum field theory. In the special limit $\eta \rightarrow 0$, the above black brane solution reduces to the well-known AdS black brane and its dual is a relativistic conformal theory. On this background geometry, we further studied the hydrodynamic transport coefficients of the vector and shear modes following the membrane paradigm [8]. In the Einstein-dilaton theory, there exists a nontrivial dilaton profile which, according to the gauge/gravity duality, corresponds to the running coupling of the dual field theory due to the non-conformality. After introducing one parameter α describing the weight of a running coupling in the dual field theory, we investigated the DC conductivity and charge diffusion constant of the general D -dimensional dual field theory. We showed that these results provide the exactly same results obtained by Kubo formula in $D = 4$ with $\alpha = 0$ or $-\eta/2$ [24].

In the previous cases [24], we simply assume the gauge/gravity duality. However, it is very interesting question to ask whether we can really apply the gauge/gravity duality to the non-AdS geometry. There were several examples using this gauge/gravity duality in the asymptotic non-AdS space, for example Lifshitz, Schrödinger geometries and Sakai-Sugimoto model, etc. Unfortunately, we do not have any exact proof for the gauge/gravity duality even in the AdS geometry, although there were many evidences for the AdS/CFT correspondence. In such situation, it is not hopeful to prove the gauge/gravity duality. Instead, we want find the possibility for the generalization of the AdS/CFT correspondence to the non-AdS geometry. That is the main goal of this work. In this paper, after assuming the gauge/gravity duality, we derived the thermodynamic properties of the dual field theory

from the black brane geometry, in which the details of the dual field theory were not used. On this thermal equilibrium, if we turn on the shear modes, their transport coefficients can be described by the Kubo formula derived from the quantum field theory and interestingly include informations of the thermal equilibrium like energy and pressure (see (63)). If the generalization to the gauge/gravity duality is possible, we can expect that the physical properties of the dual field theory obtained by the black brane thermodynamics and by the Kubo formula should be consistent. In Sec. 3.3, we studied the transports coefficient of the shear modes and calculated the sum of the energy density and pressure from the momentum diffusion constant. Furthermore, after rescaling of time or fixing the combination of two parameters Λ and ϕ_0 , we showed that the sum of the energy density and pressure in the momentum diffusion constant is exactly the same as that obtained from the black brane thermodynamics, which shows the possibility applying the AdS/CFT correspondence to the non-AdS geometry. Notice that here we used the membrane paradigm instead of the Kubo formula and checked that the membrane paradigm reproduces the same results obtained by the Kubo formula in Sec. 3.1.

Although the exact correspondence between the bulk fluctuations and their dual operators is still unclear, this work shows one possibility for the generalization of the the gauge/gravity duality to the non-AdS space. The gauge/gravity duality in the non-AdS space is one of the important issues and it is interesting to apply it to the real physical systems. We hope to report more results in near future.

Acknowledgement

This work was supported by the Korea Science and Engineering Foundation (KOSEF) grant funded by the Korea government(MEST) through the Center for Quantum Spacetime(CQUeST) of Sogang University with grant number R11-2005-021. C. Park was also supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology(2010-0022369).

References

- [1] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2** (1998) 231 [hep-th/9711200].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Phys. Lett. B* **428** (1998) 105 [hep-th/9802109].
- [3] E. Witten, *Adv. Theor. Math. Phys.* **2** (1998) 253 [hep-th/9802150].
- [4] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, *Phys. Rev. Lett.* **95**, 261602 (2005) [hep-ph/0501128]; A. Karch, E. Katz, D. T. Son and M. A. Stephanov, *Phys. Rev. D* **74**, 015005 (2006) [hep-ph/0602229]; T. Sakai and S. Sugimoto, *Prog. Theor. Phys.* **113**, 843

- (2005) [hep-th/0412141]; T. Sakai and S. Sugimoto, Prog. Theor. Phys. **114**, 1083 (2005) [hep-th/0507073]; C. P. Herzog; Phys. Rev. Lett. **98**, 091601 (2007) [hep-th/0608151]. B. -H. Lee, C. Park and S. -J. Sin, JHEP **0907**, 087 (2009) [arXiv:0905.2800 [hep-th]]; C. Park, Phys. Rev. D **81**, 045009 (2010) [arXiv:0907.0064 [hep-ph]]; K. Jo, B. -H. Lee, C. Park and S. -J. Sin, JHEP **1006**, 022 (2010) [arXiv:0909.3914 [hep-ph]]; C. Park, Phys. Lett. B **708**, 324 (2012) [arXiv:1112.0386 [hep-th]].
- [5] G. Policastro, D. T. Son and A. O. Starinets, JHEP **0209**, 043 (2002) [hep-th/0205052].
- [6] G. Policastro, D. T. Son and A. O. Starinets, JHEP **0212**, 054 (2002) [hep-th/0210220].
- [7] P. Kovtun, D. T. Son and A. O. Starinets, JHEP **0310**, 064 (2003) [hep-th/0309213].
- [8] N. Iqbal and H. Liu, Phys. Rev. D **79**, 025023 (2009) [arXiv:0809.3808 [hep-th]].
- [9] Y. Matsuo, S. -J. Sin and Y. Zhou, JHEP **1201**, 130 (2012) [arXiv:1109.2698 [hep-th]].
- [10] Y. Matsuo, S. -J. Sin and Y. Zhou, JHEP **1207**, 050 (2012) [arXiv:1204.6627 [hep-th]].
- [11] S. S. Gubser, I. R. Klebanov and A. W. Peet, Phys. Rev. D **54**, 3915 (1996) [hep-th/9602135].
- [12] S. A. Hartnoll, Class. Quant. Grav. **26**, 224002 (2009) [arXiv:0903.3246 [hep-th]]; C. P. Herzog, J. Phys. A **42**, 343001 (2009) [arXiv:0904.1975 [hep-th]]; J. McGreevy, Adv. High Energy Phys. **2010**, 723105 (2010) [arXiv:0909.0518 [hep-th]]; G. T. Horowitz, arXiv:1002.1722 [hep-th]; S. Sachdev, arXiv:1002.2947 [hep-th].
- [13] S. S. Gubser, Phys. Rev. D **78**, 065034 (2008) [arXiv:0801.2977 [hep-th]].
- [14] S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, Phys. Rev. Lett. **101**, 031601 (2008) [arXiv:0803.3295 [hep-th]].
- [15] S. S. Gubser and S. S. Pufu, JHEP **0811**, 033 (2008) [arXiv:0805.2960 [hep-th]].
- [16] S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, JHEP **0812**, 015 (2008) [arXiv:0810.1563 [hep-th]].
- [17] J. -W. Chen, Y. -J. Kao, D. Maity, W. -Y. Wen and C. -P. Yeh, Phys. Rev. D **81**, 106008 (2010) [arXiv:1003.2991 [hep-th]].
- [18] C. P. Herzog, Phys. Rev. D **81**, 126009 (2010) [arXiv:1003.3278 [hep-th]].
- [19] M. Ammon, J. Erdmenger, M. Kaminski and P. Kerner, Phys. Lett. B **680**, 516 (2009) [arXiv:0810.2316 [hep-th]].

- [20] M. Ammon, J. Erdmenger, M. Kaminski and P. Kerner, JHEP **0910**, 067 (2009) [arXiv:0903.1864 [hep-th]].
- [21] S. S. Gubser, C. P. Herzog, S. S. Pufu and T. Tesileanu, Phys. Rev. Lett. **103**, 141601 (2009) [arXiv:0907.3510 [hep-th]].
- [22] J. P. Gauntlett, J. Sonner and T. Wiseman, Phys. Rev. Lett. **103**, 151601 (2009) [arXiv:0907.3796 [hep-th]].
- [23] S. A. Hartnoll, J. Polchinski, E. Silverstein and D. Tong, “Towards strange metallic holography,” JHEP **1004**, 120 (2010) [arXiv:0912.1061 [hep-th]]; B. -H. Lee, S. Nam, D. -W. Pang and C. Park, Phys. Rev. D **83**, 066005 (2011) [arXiv:1006.0779 [hep-th]]; B. -H. Lee, D. -W. Pang and C. Park, JHEP **1007**, 057 (2010) [arXiv:1006.1719 [hep-th]].
- [24] S. Kulkarni, B. -H. Lee, C. Park and R. Roychowdhury, arXiv:1205.3883 [hep-th].
- [25] K. Goldstein, S. Kachru, S. Prakash and S. P. Trivedi, JHEP **1008**, 078 (2010) [arXiv:0911.3586 [hep-th]].
- [26] C. Charmousis, B. Gouteraux, B. S. Kim, E. Kiritsis and R. Meyer, JHEP **11**, 151 (2010) [arXiv:1005.4690 [hep-th]].
- [27] M. Cadoni, S. Mignemi and M. Serra, Phys. Rev. D **84**, 084046 (2011) [arXiv:1107.5979 [gr-qc]].
- [28] K. Goldstein, N. Iizuka, S. Kachru, S. Prakash, S. P. Trivedi and A. Westphal, JHEP **10**, 027 (2010) [arXiv:1007.2490 [hep-th]].
- [29] X. Dong, S. Harrison, S. Kachru, G. Torroba and H. Wang, arXiv:1201.1905 [hep-th].
- [30] L. Huijse, S. Sachdev and B. Swingle, Phys. Rev. B **85**, 035121 (2012) [arXiv:1112.0573 [cond-mat.str-el]].
- [31] B. S. Kim, arXiv:1202.6062 [hep-th].
- [32] E. Perlmutter, arXiv:1205.0242 [hep-th].
- [33] B. Gouteraux and E. Kiritsis, JHEP **1112**, 036 (2011) [arXiv:1107.2116 [hep-th]].
- [34] I. Kanitscheider and K. Skenderis, JHEP **0904**, 062 (2009) [arXiv:0901.1487 [hep-th]].
- [35] B. Gouteraux, J. Smolic, M. Smolic, K. Skenderis and M. Taylor, JHEP **1201**, 089 (2012) [arXiv:1110.2320 [hep-th]].
- [36] S. S. Gubser, Adv. Theor. Math. Phys. **4**, 679 (2000) [hep-th/0002160].

- [37] U. Gursoy and E. Kiritsis, JHEP **0802**, 032 (2008) [arXiv:0707.1324 [hep-th]].
- [38] U. Gursoy, E. Kiritsis and F. Nitti, JHEP **0802**, 019 (2008) [arXiv:0707.1349 [hep-th]].
- [39] B. Gouteraux and E. Kiritsis, JHEP **1112**, 036 (2011) [arXiv:1107.2116 [hep-th]].